

D-branes in $\mathcal{N} = 2$ coset models and twisted equivariant K-theory

Sakura Schäfer-Nameki [#]

*Department of Applied Mathematics and Theoretical Physics
University of Cambridge, Wilberforce Road, Cambridge CB3 0WA, U.K.*

Abstract

The charges of D-branes in Kazama-Suzuki coset models are analyzed. We provide the calculation of the corresponding twisted equivariant K-theory, and in the case of Grassmannian cosets, $su(n+1)/u(n)$, compare this to the charge lattices that are derived from boundary conformal field theory.

8.8.2003

[#] e-mail: S.Schafer-Nameki@damtp.cam.ac.uk

1. Introduction

Twisted K-theory has recently attracted much attention, both in mathematics and in string theory. Most notable among the results obtained in this subject is the theorem by Freed, Hopkins and Teleman (FHT) relating twisted K-theory to the Verlinde algebra [1,2,3], which seen in the light of efforts in string theory to relate K-theory to D-brane charges, as computed from conformal field theory data, is indeed very suggestive.

In string theory, the importance of K-theory is founded upon the conjecture that D-brane charges are classified by the K-theory of the target space [4,5]. This conjecture has been tested in various instances, when both an algebraic description of the D-branes in terms of boundary conformal field theory (BCFT) as well as the relevant K-theory are accessible to explicit computations. Most prominent examples of theories, where the BCFT is under control are free CFTs, orbifolds, WZW and coset models. The relation to K-theory has been established in a vast number of examples for the free and orbifold theories. In the case of (supersymmetric) WZW models, the presence of the non-trivial background B-field implies that D-brane charges should take values in a twisted version of K-theory. The K-theories relevant for WZW models have been computed in [6,7,8] and indicate that the known boundary states do not in general suffice to fill up the charge lattice that is predicted by K-theory.

In this paper the correspondence between D-brane charges and K-theory shall be explored for coset CFTs [9]. Supersymmetry is key to the present discussion, and the relevant models are the Kazama-Suzuki coset theories [10], of which the Grassmannian and generalized parafermionic cosets shall be of foremost interest. Using the FHT result, we provide the computation of the twisted equivariant K-groups associated to these models and discuss the relation to the BCFT.

The plan of the paper is as follows. In section 2, we discuss as a prelude the super-minimal models, and compare K-theory to BCFT in this simplest example. Section 3 provides some detail on the CFT-side of the cosets. The main result is contained in section 4, where we use the theorem by FHT [1,2] to compute the twisted equivariant K-theory of the coset models. A matching with the BCFT results is established in the case of Grassmannian cosets, and we close with concluding remarks and acknowledgements.

2. Superparafermions

The super-minimal models, realized as superparafermionic $u(1)$ cosets of the $su(2)_k$ WZW models, are the simplest examples of $\mathcal{N} = 2$ superconformal field theories. In this section we review the boundary CFT in these models, as well as compute the K-theory lattice of D-brane charges. This should serve as an appetizer to subsequent sections, where generalizations to other $\mathcal{N} = 2$ coset theories shall be discussed.

2.1. The boundary CFT side

The superparafermions [11,10] have a representation in terms of the coset models

$$\frac{su(2)_k \oplus u(1)_4}{u(1)_{2(k+2)}}. \quad (2.1)$$

The highest weight representations will be labeled by (Λ, m, a) , where Λ is an $su(2)_k$ weight, m labels the $u(1)_{2(k+2)}$ representations and a specifies the weight of $u(1)_4$. The induced level $2\kappa = 2(k+2)$ of the $u(1)_{2(k+2)}$ denominator theory indicates that the weight m takes values in $\mathbb{Z}/2(k+2)$. The theories (2.1) are obtained as supersymmetric generalizations of the coset construction for $su(2)/u(1)$, where $\dim(su(2)) - \dim(u(1))$ adjoint fermions, described by the $u(1)_4$ factor, are introduced. The corresponding representation label a takes values in $\mathbb{Z}/4$. The coset selection and identification rules are

$$\begin{aligned} \Lambda + m + a &\in 2\mathbb{Z}, \\ (\Lambda, m, a) &\equiv (k - \Lambda, m + k + 2, a + 2), \end{aligned} \quad (2.2)$$

which are obtained in the standard fashion by acting with the proper automorphism $\sigma : \Lambda = [\lambda_1] \rightarrow [k - \lambda_1]$ on the modular S-matrices [12]. The theories (2.1) are the simplest examples of the generalized parafermion cosets $su(n+1)_k/u(1)^n$, and the supersymmetric Grassmannian coset models $su(n+1)_k/u(n)$.

The boundary states for (2.1) are well known [13,14,15,16] and a brief review of the results necessary for the present discussion is in place. We shall focus on A-type boundary states for the $\mathcal{N} = 2$ algebra. In view of the comparison to K-theory, the lattice of RR-charges will be of main interest. In the BCFT this is computed by the intersection index [17], which in the case of the $\mathcal{N} = 2$ superparafermion coset models has been obtained in [13,18,15]. Recall that the intersection index in the closed string picture is defined in terms of the overlaps of the RR-sectors of the boundary states [18,15]. The key observation in [15] is that the states with $\Lambda = 0$ form a generating set for the charge lattice of boundary

states in the superparafermion models. In detail, the boundary state intersection for Cardy states $\|(\Lambda_i, m_i, 0)\rangle\rangle$ is

$$(\mathcal{I}_{(\Lambda_1, \Lambda_2)})_{m_1, m_2} = N_{\Lambda_1, \Lambda_2}^{m_1 - m_2} = \left(\sum_{\Gamma} N_{\Gamma, \Lambda_2}^{\Lambda_1} \sum_m \sum_{w \in W_{su(2)}} \epsilon_w g^{-m + m_0} \right)_{m_1, m_2}, \quad (2.3)$$

where the first sum extends over all $su(2)$ level k representations, and the second sum is over all states, that are coset fields representing Ramond sector ground states, *i.e.*, which by [19] take the form (Λ, m, a) , with $m, \Lambda \in \mathbb{Z}$, such that

$$m = w(\Lambda + 1), \quad (2.4)$$

for some element w in the $su(2)$ Weyl group, $W_{su(2)} = \langle \pm \mathbb{I} \rangle$. Further $m_0 = 1$ is the $u(1)$ charge of the spectral flow operator, and N and g are the $su(2)$ and $u(1)$ fusion matrices, respectively. In detail, the fusion matrix element for the $u(1)$ fields m, m_1, m_2 is given by $(g^m)_{m_1 m_2} = \delta_{m_1, m_2 + m}$, or equivalently, g is the shift matrix $g_{m_1 m_2} = \delta_{m_1, m_2 + 1}$. In (2.3) we have fixed $a = 0$ and view $\mathcal{I}$ as a matrix in the denominator $u(1)$ weights, m_i . By performing the sum over the $u(1)$ weights m , that are allowed by (2.4) for a given $su(2)$ weight Γ we arrive at

$$\mathcal{I}_{(\Lambda_1, \Lambda_2)} = \sum_{\Gamma} N_{\Gamma, \Lambda_2}^{\Lambda_1} (g^{-\Gamma} - g^{\Gamma+2}). \quad (2.5)$$

To prove that the $\Lambda = 0$ states, with fixed value of a , form a basis of the charge lattice, first note that

$$\mathcal{I}_{(0,0)} = \mathbb{I} - g^2. \quad (2.6)$$

The matrices (2.5) with general values for Λ_1 and Λ_2 can be obtained from the following linear combination

$$\begin{aligned} & (g^{-\Lambda_1} + g^{-\Lambda_1+2} + \dots + g^{\Lambda_1}) \mathcal{I}_{(0,0)} (g^{-\Lambda_2} + g^{-\Lambda_2+2} + \dots + g^{\Lambda_2}) \\ &= (g^{-\Lambda_1} + g^{-\Lambda_1+2} + \dots + g^{\Lambda_1}) \sum_{\Gamma} N_{\Gamma, \Lambda_2}^0 (g^{-\Gamma} - g^{\Gamma+2}) \\ &= \sum_{\Gamma} N_{\Gamma, \Lambda_2}^{\Lambda_1} (g^{-\Gamma} - g^{\Gamma+2}), \end{aligned} \quad (2.7)$$

where the last step follows by inserting the explicit form of the fusion coefficients (*e.g.* [20,21]). In particular for $su(2)$ these are

$$N_{\Gamma, \Lambda_2}^{\Lambda_1} = \sum_{\Lambda \in P_{\Lambda_1}} \sum_{w \in W_{su(2)}} \epsilon_w \delta_{w(\Lambda + \Gamma + 1) - \Lambda_2 - 1}, \quad (2.8)$$

where P_Λ is the weight system of Λ . Thus all boundary intersections can be written in terms of the intersection of $\Lambda = 0$ states. (2.6) and the explicit form of the matrix g , which shifts every $u(1)$ label by one, implies that the rank of $\mathcal{I}_{(0,0)}$ is $k + 1$.

There are two types of A-branes, which satisfy the gluing conditions $G^\pm = \pm \bar{G}^\mp$, which were referred to as even (odd) branes in [14]. Naively, each of these give rise to a charge lattice $\mathbb{Z}^{k+1}$, thus resulting in the total charge lattice $\mathbb{Z}^{2(k+1)}$, which is isomorphic to the Verlinde fusion ring of $u(1)_{2(k+1)}$, *i.e.*

$$V_{u(1)_{2(k+1)}} = \frac{R_{u(1)}}{I_{2(k+1)}} = \mathbb{Z}^{2(k+1)}, \quad (2.9)$$

where $R_{u(1)} \cong \mathbb{Z}[\zeta, \zeta^{-1}]$ is the representation ring of $u(1)$ and $I_{2(k+1)}$ is the Verlinde ideal.

Taking the point of view of the gauged WZW model, the $\Lambda = 0$ states can be seen to form a basis for the charge lattice as follows. As explained in [22,14], the parafermions (2.1) have a disk target space (together with a non-trivial B-field and dilaton). For fixed level k we introduce $k + 2$ even and $k + 2$ odd points on the boundary of the disk. Then it was shown in [14] that there are $(k+2)(k+1)$ even (odd) A-branes, corresponding to D1-branes stretching between the even (odd) points. The $\Lambda = 0$ D1-branes $\|0, n, s\rangle\rangle$ stretch between consecutive points (of same chirality, that is, even or odd), *i.e.*, between $(n-1)\pi/(k+2)$ and $(n+1)\pi/(k+2)$, and are thus the shortest D1-branes. From the tachyons analysis in [14] we follow that two consecutive such D1-branes $\|0, n, s\rangle\rangle$ and $\|0, n+2, s\rangle\rangle$, decay and merge to give a D1-brane with non-trivial Λ , $\|1, n+1, s\rangle\rangle$, as depicted in Figure 1. Iterating this process we can produce all branes (for fixed chirality) from the $\Lambda = 0$ states, as expected from the above intersection index computation. These decays have been analyzed also in [23,16].




Figure 1 Decay of two $\Lambda = 0$ into a $\Lambda = 1$ D-brane for $k = 6$.

A simple counting argument then provides the charge lattice: there are $k + 2$ even points, and therefore $k + 2$ D1-branes with $\Lambda = 0$. Since in addition the full ring of $\Lambda = 0$ states is trivial, one of these branes is redundant. The charge lattice for the even branes is therefore $\mathbb{Z}^{k+1}$. The same argument holds for the odd branes, so that the total charge lattice is $\mathbb{Z}^{2(k+1)}$. This discussion is therefore in agreement with the calculation in terms of the intersection index.

2.2. Twisted equivariant K-theory ${}^\tau K_{U(1)}(SU(2))$

This section provides a comparison of the D-brane charges, as computed in the last section, with K-theory. The gauged WZW description dictates what the relevant K-theory for these backgrounds is. The gauging implies that the K-theory needs to be equivariant with respect to the conjugation action, and the presence of the non-trivial B-field necessitates a non-trivial twisting with respect to a 3-form. For other discussion of equivariant and twisted K-theory see *e.g.* [24,25,26,27,28,31,29].

In the case of the $SU(2)/U(1)$ coset we can compute the twisted equivariant K-group directly, using a Mayer-Vietoris argument. This calculation generalizes in principle to $SU(N)$ with $N > 2$, but the story gets more involved due to the rather complicated cell structure (see *e.g.* [30]) and other ways of computation will be more suitable. A calculation similar in spirit to the one presented in this section can be found in [1] for ${}^\tau K_{SU(2)}(SU(2))$.

The geometrical action corresponding to the CFT coset is the conjugation action. Thus, the action of $U(1)$ on $SU(2)$ is

$$SU(2) \ni g \mapsto \text{diag}(e^{it}, e^{-it}) g \text{diag}(e^{-it}, e^{it}). \quad (2.10)$$

The fixed point set is an S^1 . We choose an open covering of $SU(2) \sim S^3$ that is invariant under this action

$$A = SU(2) \setminus \{-id\}, \quad B = SU(2) \setminus \{id\}, \quad (2.11)$$

where A and B intersect in the equatorial S^2 . The Mayer-Vietoris sequence for this CW triad then reads

$$\begin{array}{ccccccc} {}^\tau K_{U(1)}^0(SU(2)) & \longrightarrow & {}^\tau K_{U(1)}^0(A) \oplus {}^\tau K_{U(1)}^0(B) & \xrightarrow{\varphi} & {}^\tau K_{U(1)}^0(A \cap B) \\ \uparrow & & & & \downarrow \\ {}^\tau K_{U(1)}^1(A \cap B) & \longleftarrow & {}^\tau K_{U(1)}^1(A) \oplus {}^\tau K_{U(1)}^1(B) & \longleftarrow & {}^\tau K_{U(1)}^1(SU(2)) \end{array} \quad (2.12)$$

Since the fixed S^1 intersects A as well as B in a point, the corresponding groups are simply given by the representation ring of $U(1)$, *i.e.*

$$\tau K_{U(1)}^0(A) = \tau K_{U(1)}^0(B) = R_{U(1)}, \quad \tau K_{U(1)}^1(A) = \tau K_{U(1)}^1(B) = 0. \quad (2.13)$$

To compute $\tau K_{U(1)}^*(A \cap B)$ we need to apply a second Mayer-Vietoris argument. So, choosing the analogous covering for $A \cap B = S^2$ as in (2.11), we obtain

$$\begin{array}{ccccc} \tau K_{U(1)}^0(S^2) & \longrightarrow & R_{U(1)} \oplus R_{U(1)} & \xrightarrow{\psi} & \tau K_{U(1)}^0(S^1) \\ \uparrow & & & & \downarrow \\ 0 & \longleftarrow & 0 & \longleftarrow & \tau K_{U(1)}^1(S^2) \end{array}. \quad (2.14)$$

Now note that $S^1 = U(1)/\mathbb{Z}_2$. To see this, note that the S^1 in the intersection of the covers of S^2 is embedded into the $SU(2)$ by

$$S^1 \cong \left\langle \begin{pmatrix} 0 & e^{i\theta} \\ -e^{-i\theta} & 0 \end{pmatrix} \right\rangle, \quad (2.15)$$

whereupon the action of the $U(1)$ as of (2.10) is by twice the angle. Thus

$$\tau K_{U(1)}^0(S^1) = \tau K_{\mathbb{Z}_2}^0(pt.) = \mathbb{Z}^2, \quad (2.16)$$

the map ψ is given by

$$\psi : (\chi, \chi') \mapsto (\chi(1) - \chi'(1), \chi(-1) - \chi'(-1)). \quad (2.17)$$

Hence the cokernel is $\tau K_{U(1)}^1(A \cap B) = 0$ and the kernel is

$$\tau K_{U(1)}^0(A \cap B) = (R_{U(1)} \oplus R_{U(1)})/\psi, \quad (2.18)$$

i.e., these are pairs of characters identified at id and $-id$, and so the quotient by ψ removes two copies of $\mathbb{Z}$.

Putting this together, the map φ in (2.12) becomes

$$\begin{aligned} \varphi : R_{U(1)} \oplus R_{U(1)} &\rightarrow (R_{U(1)} \oplus R_{U(1)}) \\ (f, g) &\mapsto (f + gL^\kappa, f + gL^{-\kappa}), \end{aligned} \quad (2.19)$$

where L is the generator of $R_{U(1)}$ and we have used the fact that due to the twisting, the bundles over A and B are patched together by tensoring with the twist-class. So this has trivial kernel, *i.e.*, $\tau K_{U(1)}^0(SU(2)) = 0$. The cokernel, (without the ψ identification) would

be $(R_{U(1)} \oplus R_{U(1)})/I_\kappa$, where the ideal is $I_\kappa = (L^\kappa - L^{-\kappa})$, and so the cokernel would be isomorphic to $\mathbb{Z}^{2\kappa}$. Including the ψ identification then results in $\mathbb{Z}^{2\kappa-2}$. We conclude thereby that

$$\tau K_{U(1)}^1(SU(2)) = \mathbb{Z}^{2\kappa-2} = \mathbb{Z}^{2(k+1)}, \quad \tau K_{U(1)}^0(SU(2)) = 0. \quad (2.20)$$

Comparison with (2.9) then yields that

$$K_{U(1)}^1(SU(2)) = \frac{R_{u(1)}}{I_{2(k+1)}}, \quad (2.21)$$

which is reminiscent of the statement in [1,3] in the case of the topological G/G coset and G -equivariant twisted K-theory of G .

2.3. Comparison between K-theory and D-brane charges

In the comparison between K-theory and D-brane charges there are two subtle points that one has to take into account. Firstly, in order to compare the K-theory to the CFT, we note, that not only the equivariance with respect to the $u(1)$ action has to be taken into account. Recall that the selection and identification rules amount to modding out in addition with the common centre of $su(2)$ and $u(1)$, *i.e.*, with $\mathbb{Z}_2$ as specified in the second equation of (2.2). Hence, the CFT needs to be compared to $\tau K_{U(1)/\mathbb{Z}_2}(SU(2))$. The key change that occurs in the above K-theory computation is then that (2.16) is replaced by

$$\tau K_{U(1)/\mathbb{Z}_2}^0(S^1) = \tau K^0(pt.) = \mathbb{Z}, \quad (2.22)$$

wherefore

$$\tau K_{U(1)/\mathbb{Z}_2}^1(SU(2)) = \mathbb{Z}^{\kappa-1} = \mathbb{Z}^{k+1}, \quad \tau K_{U(1)/\mathbb{Z}_2}^0(SU(2)) = 0. \quad (2.23)$$

This corresponds precisely to the charges carried by the even (or odd) branes.

The second point to note is that K-theory is meant to classify the charges of D-branes in string theory (as opposed to CFT). In addition to conformal field-theoretical consistency conditions, in string theory the GSO-projection has to be imposed upon the boundary states. As is familiar from other super-string backgrounds (see *e.g.*, [31]), this does indeed cut down the number of allowed D-branes. *E.g.* in flat space Type II theories, GSO-invariance of the boundary states implies that both the NSNS and the RR sector parts of the boundary states have to be GSO-invariant, which implies that there are only half as many GSO-invariant boundary states than Ishibashi states (as opposed to BCFT, where the number of Cardy states equals the number of Ishibashi states). We conjecture that a similar reduction occurs for the above coset models. In the present case the NS (R) sector corresponds to even (odd) a , *i.e.*, the NSNS and RR sector branes are precisely the even and odd branes, respectively. To confirm this, one should consider a proper, critical string background based on these coset theories, such as are provided by Gepner models.

3. $\mathcal{N} = 2$ coset models and D-brane charges

In the following, we shall generalize the computations from $SU(2)/U(1)$ to the Grassmann models and generalized parafermions. We will step back for a moment and provide some relevant facts of these $\mathcal{N} = 2$ coset CFTs.

3.1. Supersymmetric coset models

Recall that any WZW model for a compact Lie group G , described by affine $\mathfrak{g}_k$ at level k , can be supersymmetrized to an $\mathcal{N} = 1$ SCFT by introducing $\dim(\mathfrak{g})$ fermionic fields transforming in the adjoint representation [32]. This theory is most elegantly described by ‘decoupling’ the bosonic and fermionic degrees of freedom (see *e.g.* [10]), which results in shifting the level of the KM theory. The $\mathcal{N} = 1$ supersymmetric version for $\mathfrak{g}_k$ is then

$$\mathfrak{g}_{k-g} \oplus so(\dim(\mathfrak{g}))_1, \quad (3.1)$$

where g is the dual Coxeter number such that $f_{acd}f_{bcd} = g\delta_{ab}$. The central charge of the model is thus

$$c = \frac{1}{2} \dim(\mathfrak{g}) + \frac{(k-g) \dim(\mathfrak{g})}{k}. \quad (3.2)$$

Due to the product structure of (3.1), it usually suffices to study the bosonic $\mathfrak{g}_k$ model and infer from there the supersymmetric statements by tensoring with the $so(d)_1$ theory. In particular, this was used in [6] to compare the boundary states to K-theory charges.

The supersymmetrization of coset models is slightly more intricate. Let $\mathfrak{h}$ be a sub-Lie algebra of $\mathfrak{g}$. The supersymmetric construction for $\mathfrak{g}_k$ contains one for $\mathfrak{h}$, and one can apply the coset construction to this pair of theories. The so-obtained $\mathcal{N} = 1$ supersymmetric coset models are of the form

$$\frac{\mathfrak{g}_k \oplus so(2d)_1}{\mathfrak{h}_{k+g_{\mathfrak{g}}-g_{\mathfrak{h}}}}, \quad (3.3)$$

where $d = \frac{1}{2}(\dim(\mathfrak{g}) - \dim(\mathfrak{h}))$ and g denotes the dual Coxeter numbers of the respective Lie algebras. The $so(2d)_1$ part again parametrizes d complex fermions. Note that this corresponds to the supersymmetric version of $\mathfrak{g}_{k+g_{\mathfrak{g}}}/\mathfrak{h}$. It is important to realize that $\mathfrak{h}$ is embedded diagonally into $\mathfrak{g}_k \oplus so(2d)_1$ and hence, that the theory does not simply factorize into bosonic and fermionic contributions, as *e.g.* in the WZW models. This is apparent from the way the $\mathfrak{h}$ current algebra is constructed in [10]. One consequence of this for the present purposes is that the bosonic results for coset branes in $\mathfrak{g}_k/\mathfrak{h}$ will not simply carry over to the supersymmetrized theory. There are non-trivial selection rules

for highest weights in the supersymmetric coset, which non-trivially mix the weights of $\mathfrak{g}$, $so(2d)$ and $\mathfrak{h}$ (see below).

Generically, (3.3) is an $\mathcal{N} = 1$ theory. As is well known from [10], for $\text{rank}(\mathfrak{g}) = \text{rank}(\mathfrak{h})$, the coset theory is $\mathcal{N} = 2$ if the geometric (right-action) coset space G/H of the corresponding Lie groups is Kähler. The representations of the coset model (3.3) are labeled by highest weights (Λ, λ, a) , which correspond to $\mathfrak{g}_k$, $\mathfrak{h}_{k+g_{\mathfrak{g}}-g_{\mathfrak{h}}}$ and $so(2d)_1$ highest weights, respectively. However, not all combinations of weights are distinct in the coset theory. Gepner developed a general method how to extract the field identifications in coset models by analyzing the action of certain automorphisms on the S-modular matrix [12].

3.2. Grassmannian coset models $Gr(n, k)$

The models of interest in the following are the generalized super-conformal minimal models based on the Grassmannians $G(n, n+k)$ of n -planes in $\mathbb{C}^{n+k}$, which after level-rank duality take the form of projective cosets

$$Gr(n, k) : \frac{su(n+1)_k \oplus so(2n)_1}{su(n)_{k+1} \oplus u(1)_{n(n+1)(k+n+1)}}. \quad (3.4)$$

The level of $u(1)_{n(n+1)(k+n+1)}$ indicates the weights are $\mathbb{Z}_{n(n+1)(k+n+1)}$ -valued. The central charge is $c = 3nk/(k+n+1)$ and the weights are labeled by (Λ, λ, m, a) , where λ and m correspond to $su(n)$ and $u(1)$ weights, respectively, as well as $so(2n)_1$ weights $a \in \mathbb{Z}/4$. The highest weights in the basis of fundamental weights are

$$\Lambda = (\Lambda_1, \dots, \Lambda_n), \quad \lambda = (\Lambda_1, \dots, \Lambda_{n-1}), \quad m = (n+1)\Lambda^{(n)} \cdot \Lambda = \sum_{i=1}^n i\Lambda_i. \quad (3.5)$$

Not all combinations of weights in (3.5) are distinct in the theory and certain field identifications need to be imposed. Recall from [12] that an automorphism σ of the extended Dynkin diagram of $\mathfrak{g}_k$ is defined to be *proper*, if

$$\sigma(\lambda) = \sigma(0) + w_{\sigma}(\lambda), \quad (3.6)$$

for some fixed $w_{\sigma} \in W$. For any proper automorphism $\sigma(0) = k\lambda_{\sigma}$, where λ_{σ} is a level 1 weight (specified for each such σ). The field identification rules are extracted by considering the action of proper automorphisms on the S-modular matrix. Let $S_{\Lambda, \Gamma}^{\mathfrak{g}, k}$ be the S-modular matrix for $\mathfrak{g}_k$. Using the explicit form of S it follows that

$$S_{\sigma(\Lambda), \Gamma}^{\mathfrak{g}, k} = e^{2\pi i \lambda_{\sigma} \Gamma} S_{\Lambda, \Gamma}^{\mathfrak{g}, k}. \quad (3.7)$$

Then, fields for which the phase in (3.7) vanishes need to be identified in order to ensure that the modular matrix has full rank.

Consider thus the proper automorphisms, acting on $su(n+1)$ and $so(2d)_1$, respectively, as

$$\sigma(\Lambda) = k\Lambda^{(1)} + w_\sigma(\Lambda), \quad \sigma_v(a) = v + w_{\sigma_v}(a), \quad (3.8)$$

where v is the $so(2d)_1$ vector weight. Note that for $su(n+1)$ the proper outer automorphisms are given by the order $n+1$ rotation group of the extended Dynkin diagram, *i.e.*

$$\sigma \left(\sum_{i=1}^n \lambda_i \Lambda^{(i)} \right) = (k - \lambda_1 - \dots - \lambda_n) \Lambda^{(1)} + \lambda_1 \Lambda^{(2)} + \dots + \lambda_{n-1} \Lambda^{(n)}. \quad (3.9)$$

There are two combinations of these automorphisms that can be applied to the coset fields

$$\begin{aligned} J_1(\Lambda, \lambda, m, a) &= (\sigma(\Lambda), \sigma(\lambda), m + k + n + 1, \sigma_v(a)), \\ J_2(\Lambda, \lambda, m, a) &= (\sigma(\Lambda), \lambda, m + n(k + n + 1), \sigma_v^n(a)), \end{aligned} \quad (3.10)$$

which are of order $n(n+1)$ and $n+1$ respectively, since σ on $su(n)$ is of order n . Applying (3.7) and (3.10), shows that the phase vanishes, and thus that the S-matrix elements for the coset field and its images under J_1 and J_2 are the same (*cf.* [19,12]). This leads us to identify the fields (3.10) in order to maintain unitarity of the S-matrix. For later reference, recall that the dimension of the chiral ring is

$$\dim \mathcal{R}^{Gr(n,k)} = \frac{(n+k)!}{n!k!}.$$

A peculiar property of the $Gr(n, k)$ is that they are conjectured to be invariant under the exchange of k and n , *i.e.*,

$$Gr(n, k) \cong Gr(k, n). \quad (3.11)$$

Despite the lack of a full proof of this level-rank duality, very convincing arguments have been put forward, most notably in [10], where it is shown that the coset representation of the $\mathcal{N} = 2$ superconformal algebra is symmetric in k and n . Furthermore, in [33] an isomorphism between the chiral data is constructed. This implies in particular, that the corresponding boundary theories need to respect level-rank duality.

The boundary intersection index $\mathcal{I}_{ab} = \text{Tr}_{\mathcal{H}_{ab}^{open}}(-1)^F = {}_{RR}\langle\langle a||b \rangle\rangle_{RR}$ [17,18,13] for the Grassmannian coset models has been computed in [15]. It was indicated there that, as discussed earlier for the superminimal model, it is sufficient to compute the index for the

$\Lambda = 0$ Cardy boundary states. The rank of the charge lattice is equal to the dimension of the chiral ring, *i.e.*,

$$\binom{n+k}{n}. \quad (3.12)$$

The result (3.12) refers to the charges of even or odd maximally supersymmetric D-branes in the Grassmannian coset CFTs and the purpose of the subsequent section is to derive this result for the charge lattice by a K-theory computation.

3.3. Superparafermion coset models $SPF(n, k)$

A related coset model is that of $SU(n)$ by its maximal torus, *i.e.* the generalized superparafermions,

$$SPF(n, k) : \frac{su(n+1)_k \oplus so(n(n+1))_1}{u(1)^n}. \quad (3.13)$$

The BCFT will be discussed elsewhere, however, we shall provide the corresponding K-theory calculation below. The selection and identification rules which are needed for this computation are as follows.

The center of $SU(n+1)$, or equivalently the outer automorphism group of its affine version, is generated by the order $n+1$ rotation σ of the extended Dynkin diagram. Again, σ_v acts on the $so(n(n+1))_1$, with $\sigma_v(0) = v$. This imposes the identification rule

$$(\sigma(\Lambda), \mathbf{m} + \sigma(0), \sigma_v^n(a)) = (\Lambda, \mathbf{m}, a), \quad (3.14)$$

where $\sigma(0) = (k+n+1)\Lambda^{(1)}$.

The dimension of the chiral ring in this case, using [19], is computed to be

$$\dim \mathcal{R}^{SPF(n,k)} = \frac{(k+n)!}{k!}.$$

4. Twisted equivariant K-theory

This section provides the computation of the twisted equivariant K-groups relevant for the generalized parafermions and Grassmannian coset models.

4.1. ${}^\tau K_H(G)$

Let G be a simple, simply-connected, connected Lie group (*i.e.*, in particular π_1 is torsion-free). For most purposes we shall consider $G = SU(N+1)$. Further let H be a connected, maximal rank subgroup of G . Then the computation of ${}^\tau K_H(G)$ can be traced back to the computation of Freed, Hopkins and Teleman in [1,2] of ${}^\tau K_G(G)$. The theorem by FHT states, that twisted G -equivariant K-theory of G is isomorphic as an algebra to the Verlinde algebra of G at a level k determined by the 3-form that specifies the twist τ of the K-theory

$${}^\tau K_G^{\dim(G)}(G) = V_k(G). \quad (4.1)$$

The twist τ given by the $k + g$ multiple of the generator of $H^3(G, \mathbb{Z})$, which corresponds to the $\mathbf{H} = dB$ 3-form flux. The important point about (4.1) that stresses its possible relevance for coset models is that the G -equivariance is taken with respect to the adjoint action of G on itself.

The case of coset models, where H is a proper, connected, maximal rank subgroup, relates to this result as follows. First define

$$G^c \times_H G = \{(g, \tilde{g}) \in G \times G; (g, \tilde{g}) \sim (h^{-1}gh, \tilde{g}h), h \in H\}. \quad (4.2)$$

Here, G^c indicates that the action of H on G is taken to be the conjugation action, whereas on the second factor, the action is the right action. Then the map

$$\begin{aligned} \Phi : \quad G^c \times_H G &\rightarrow G^c \times G/H \\ (g, \tilde{g}) &\mapsto (\tilde{g}g\tilde{g}^{-1}, \tilde{g}H), \end{aligned} \quad (4.3)$$

is well-defined and an isomorphism. In order to calculate the twisted equivariant K-groups, note further, that for any H -space

$$K_G(X \times_H G) \cong K_H(X), \quad (4.4)$$

where the equivariance is with respect to the specified action of H on X . This is simply the statement that any G -bundle over $G \times_H X$ is determined by the underlying H -bundle over X . Hence, setting $G^c = X$ we obtain

$$K_G(G^c \times G/H) = K_G(G^c \times_H G) \cong K_H(G^c), \quad (4.5)$$

For $\pi_1(G)$ torsion-free, an equivariant form of the Künneth formula holds by a result of Hodgkin and Snaith [34,35], whereby^b

$${}^\tau K_H(G) = {}^\tau K_G(G) \otimes_{R_G} R_H, \quad (4.6)$$

where R denotes as before the representation rings of the respective groups. Note that we used the isomorphism $K_G(G/H) \cong K_H(pt.)$, which is true since any G -bundle on G/H is of the form $G \times_H R \rightarrow G \times_H pt. = G/H$ for a H -representation R .

We can now complete our computation of $K_H(G)$. For R_H free as an R_G -module, which holds for H of maximal rank and connected [36], we can use the FHT result (4.1) to simplify (4.6) into

$${}^\tau K_H(G) = \frac{R_H}{I_G}. \quad (4.7)$$

I_G is the Verlinde ideal of G , that is, the ideal, that provides the identifications necessary in order to obtain the Verlinde algebra at a specified level from the representation ring of G as $V^G = R_G/I_G$. The Schubert-like combinatorial relations, that define the ideal I_G were obtained in [37].

First, in order to elucidate (4.7), it is useful to recall how the representation rings of G and H are related. Since any G -representation can be decomposed into H -representations, $R_G < R_H$. More precisely, by definition the representation ring is

$$R_G \cong \mathbb{Z}[M^*]^{W_G} \cong \mathbb{Z}[\Lambda^{(1)}, \dots, \Lambda^{(n)}], \quad (4.8)$$

where M^* is the weight lattice, W_G the Weyl group, so that $\mathbb{Z}[M^*]^{W_G}$ is the polynomial ring of invariant polynomials in the weights. Since any representation can be decomposed into representations with highest weights equal to the fundamental weights $\Lambda^{(i)}$, the latter form a basis of the representation ring.

^b More precisely, the theorem states the existence of a Künneth spectral sequence, with $E_2^{-p,q} = \text{Tor}_{R_G}^p({}^\tau K_G^q(G), K_H^q(pt.))$ converging to ${}^\tau K_H(G)$. However, in the present context R_H is a free module over R_G , wherefore the higher Tor's vanish and the spectral sequence collapses. See *e.g.* [8] for a case where this spectral sequence is non-trivial.

4.2. Twist-class and level of the Verlinde-ideal

To compare the above results with the intersection index calculation it is interesting to compute the induced level of the Verlinde ideal entering the K-theory calculation in (4.7). Recall, that each invariant symmetric bilinear form on the Lie algebra $\mathfrak{g}$ is in one to one correspondence with central extensions $\mathbb{R} \rightarrow \widetilde{L\mathfrak{g}} \rightarrow L\mathfrak{g}$ of the loop algebra [38]. This relation follows by noting that each such bilinear form $\langle, \rangle$ on $\mathfrak{g}$ defines a cocycle $\omega : L\mathfrak{g} \times L\mathfrak{g} \rightarrow \mathbb{R}$ on the corresponding loop algebra $L\mathfrak{g}$ as

$$\omega(\xi(z), \eta(z)) = \text{Res}_{z=0} \langle \xi(z), d\eta(z) \rangle. \quad (4.9)$$

If ω is integral then there is a corresponding central extension $U(1) \rightarrow \widetilde{LG} \rightarrow LG$ of the loop group LG .

In the case of ${}^\tau K_G(G)$, the twist-class is an equivariant integral cohomology class τ in $H_G^3(G)$. In [3] the twistings were chosen, such that the restriction to $H_T^3(T)$, where T is the maximal torus of G , have trivial components in $H^3(T)$. In this case it was shown that the twisted equivariant K-theory with complex coefficients can be computed as a twisted cohomology with coefficients in a line-bundle $\mathcal{L}^\tau$, which is determined by the class τ ,

$${}^\tau K_G^{0/1}(G; \mathbb{C})_g^\wedge = {}^\tau H^{even/odd}(G^g; \mathcal{L}^\tau(g)), \quad (4.10)$$

where $^\wedge$ denotes completion (see *e.g.* [39]). $\mathcal{L}^\tau$ is determined by τ as follows. $\tau \in H^2(BT) \otimes H^1(T)$, by the assumption that $\tau|_{H^3(T)} = 0$. Thus, it defines a map $\tau : T \rightarrow T^\vee$. The sheaf $\mathcal{L}^\tau(t)$ for $t \in T$ is $\tau(t)$, which specifies a line bundle over T , since $T^\vee$ labels line-bundles over T . ${}^\tau K_G^{0/1}(G; \mathbb{C})_g^\wedge$ can be viewed as the completion of sections of a sheaf (denoted ${}^\tau \mathcal{K}$ in [3]) over the GIT-quotient $G_{\mathbb{C}} // G_{\mathbb{C}}$. The simplification that occurs in the above $K_G(G)$ case is that ${}^\tau \mathcal{K}$ is a skyscraper-sheaf with one-dimensional stalks, and thus the twisted K-theory can be calculated exactly by means of this completion technique. Note, that these results hold for K-theory with complex coefficients, and thus in particular do not take into account torsion. The case of integral K-theory will be discussed in [40].

All this extends to the case of interest in this paper, *i.e.*, H -equivariant K-theory of G . The twistings are in $H_H^3(G)$, and we restrict them to $H_{T_H}^3(T_G)$, whereby T_G is the maximal torus of G . Again, in $H_{T_H}^3(T_G) = H^3(ET_H \times_{T_H} T_G)$ we assume that τ restricts trivially to $H^3(T_H)$. Thus τ gives rise to a map $T_H \rightarrow T_H^\vee$, thus specifying a line-bundle over each point in T_H .

A comment on induced level. Fix an element $\tau \in H_H^3(G)$ to be the (topological) level of a central extension of LG . Then we can write this as $k + g$ times the generator of $H_H^3(G)$, with k the level and g the dual Coxeter number. The induced twisting on a subgroup H will be $k + g$, wherefore the induced level is $k + g - g_H$, where g_H is the dual Coxeter number for H .

4.3. $K_{U(1)}(SU(2))$ revisited

As a first application of the above results, we revisit the calculation for $SU(2)/U(1)$, which we already performed using the cell complex earlier. The representation ring of $SU(2)$ is $R_{SU(2)} = \mathbb{Z}[\Lambda]$, where Λ labels the fundamental representation. The ring $R_{U(1)}$ is generated by the one-dimensional representation. More precisely, a representation of $U(1)$ is specified by an integer p , such that

$$\begin{aligned} \zeta^p : U(1) &\rightarrow GL(\mathbb{C}) \\ e^{i\theta} &\mapsto e^{ip\theta}. \end{aligned} \quad , \quad (4.11)$$

i.e., $U(1)$ acts by multiplication with the p th powers of its generators. Since $\zeta^p \otimes \zeta^q = \zeta^{p+q}$, the representation ring is generated by ζ, ζ^{-1} and is isomorphic to $\mathbb{Z}[\zeta, \zeta^{-1}]$. The twisting for $SU(2)$ is $k+2$ and the standard Verlinde ideal is[#] $I_{SU(2)}^k = \langle \Lambda^{k+1} \rangle$. To compute (4.7) in this context we note that the representation ring of $SU(2)$ is embedded into that of $U(1)$ by the identification $\Lambda = \zeta + \zeta^{-1}$, and thus

$${}^\tau K_{U(1)}(SU(2)) = \frac{R_{SU(2)}}{\langle \Lambda^{k+1} \rangle} \otimes R_{U(1)} = \frac{\mathbb{Z}[\zeta]}{\langle (\zeta + \zeta^{-1})^{k+1} \rangle} = \mathbb{Z}^{2(k+1)}, \quad (4.12)$$

in degree $\dim(SU(2))$, *i.e.* for all odd degrees, and vanishes otherwise. This is precisely our result (2.20). The induced level for the Verlinde ideal as embedded into $R_{U(1)}$ is again k and the twisting $\tau = (k+2)$.

4.4. $K_{U(n)}(SU(n+1))$

Next we generalize this to the Grassmannian coset models. First recall that

$$V_{SU(n+1)}^k \cong \mathbb{Z}^{\binom{n+k}{k}}. \quad (4.13)$$

The twisting in $SU(n)$ is $k+n+1$ times the generator of $H_H^3(G)$. By the reasoning in the last section, the twisting for the $SU(n)$ subgroup is thus also $k+n+1$, which corresponds to the level $k+1$ in $SU(n)$. (4.7) implies that

$${}^\tau K_{U(n)}(SU(n+1)) = \frac{R_{U(n)}}{I_{SU(n+1)}^k}, \quad (4.14)$$

[#] In the following, only the $\mathbb{Z}$ -module structure is taken into account, since our main interest is the rank of the charge lattice, rather than the ring structure.

in degree $\dim(SU(n+1))$ and vanishes for the complementary degrees. This can be made more explicit as follows. From Theorem 3 in [36] we know that for a maximal rank subgroup, which is connected and closed

$$K(G/H) \cong R_H \otimes_{R_G} \mathbb{Z}, \quad (4.15)$$

where G/H is the right quotient. Applied to the present situation, where $SU(n+1)/U(n) = \mathbb{C}P^n$, and using the K-theory for complex projective space as computed by Adams in [41], this reads

$$R_H \otimes_{R_G} \mathbb{Z} \cong K(\mathbb{C}P^n) \cong \mathbb{Z}^{n+1}. \quad (4.16)$$

Together with (4.13), (4.14) simplifies therefore to

$${}^\tau K_{U(n)}(SU(n+1)) \cong \mathbb{Z}^{(n+1) \binom{n+k}{k}}. \quad (4.17)$$

4.5. $K_{U(1)^n}(SU(n+1))$

For generalized super-parafermion cosets, we invoke Theorem 1 in [36], which implies, that $R_{u(1)^n}$ is free of rank $(n+1)!$, *i.e.*,

$$R_{U(1)^n} \cong (R_{SU(n+1)})^{|W|} = (R_{SU(n+1)})^{(n+1)!}, \quad (4.18)$$

where $|W|$ is the order of the Weyl group. Therefore (4.7) results in

$${}^\tau K_{U(1)^n}(SU(n+1)) = \left(\frac{R_{SU(n+1)}}{I_{SU(n+1)}^k} \right)^{\oplus (n+1)!} = \mathbb{Z}^{(n+1)! \binom{n+k}{k}}. \quad (4.19)$$

4.6. Comparison between K-theory and D-brane charges

To compare the above K-theoretical results to the charge lattice that the boundary conformal field theory and string theory predicts, there are again two subtle points that need to be elucidated. Firstly, as in the $SU(2)/U(1)$ case, the identification rules implied by the order $n(n+1)$ outer automorphism (3.9) necessitate that the K-theory has to be computed with respect to an additional equivariance. The automorphisms (3.10) act on the affine $su(n+1)$ weights as outer automorphisms, and thus correspond to central elements of the corresponding Lie group $SU(n+1)$. For this reason, the equivariance in the K-theory for $Gr(n, k)$ is with respect to $U(n)/Z$, where

$$Z = \mathbb{Z}_{n(n+1)} = \left\langle e^{2\pi i/(n+1)} \mathbb{I}_{n+1}, \begin{pmatrix} e^{2\pi i/n} \mathbb{I}_n & 0 \\ 0 & 1 \end{pmatrix} \right\rangle, \quad (4.20)$$

where the first element generates the center of $SU(n+1)$ and the second generates the center of $SU(n)$. This implies

$$H/Z = U(n)/\mathbb{Z}_{n(n+1)} = \frac{SU(n) \times U(1)}{\mathbb{Z}_n} \Big/ \mathbb{Z}_{n(n+1)} = U(n)/\mathbb{Z}_{n+1}. \quad (4.21)$$

Therefore there is an additional $\mathbb{Z}_{n+1}$ identification that needs to be taken into account in the K-theory computation, in order to compare with the boundary conformal field theory. Now note that $\mathbb{Z}_{n+1}$ sits inside $U(1) \subset U(n)$, so that the condition on the subgroup being connected still holds. That the above arguments goes through for the H/Z -equivariant K-theory[‡], is seen most directly in the formulation of [3]. Recall from section 4.2, that the twisted equivariant K-theory ${}^\tau K_H(G)$ can be computed by taking sections of a sheaf ${}^\tau \mathcal{K}(G)$, which is defined on $H_{\mathbb{C}}//H_{\mathbb{C}}$. There is a natural Z -action on $H_{\mathbb{C}}//H_{\mathbb{C}}$, and in order to compute the H/Z -equivariant K-theory, one again takes sections of ${}^\tau \mathcal{K}(G)$, whose stalks are still 1-dimensional, however, the Z -action forces to take invariant sections, which amounts to identifying the stalks over the points in each orbit of Z . This leads for the present case to an $(n+1)$ -fold identification of the stalks, and thus, the K-groups, *i.e.*, the sections of the sheaf in question, are computed to be

$${}^\tau K_{U(n)/\mathbb{Z}_{n+1}}(SU(n+1)) = \mathbb{Z}^{\binom{n+k}{k}}. \quad (4.22)$$

Note that this is in complete agreement with the charges obtained from the even (odd) Cardy branes in the CFT. Further, it is consistent with level-rank duality.

Analogously, for the generalized superparafermion cosets, the automorphism is of order $n+1$ and similar reasoning implies that the charge lattice is again given by the dimension of the chiral ring

$${}^\tau K_{U(1)^n/\mathbb{Z}_{n+1}}(SU(n+1)) = \mathbb{Z}^{n! \binom{n+k}{k}}. \quad (4.23)$$

It would be interesting to check this on the CFT side.

The second point that needs to be addressed refers to the D-brane charges as one would expect them from string theory. As discussed in the case of $SU(2)/U(1)$, a full string theoretical treatment would require to GSO project, which in particular necessitates to cut down the D-brane spectrum to particular superpositions of RR and NSNS sector boundary states. This may serve as an explanation, why the K-theory sees only half of the brane charges of the CFT.

[‡] I thank S. Fredenhagen for pointing out some subtleties in this argument.

5. Conclusions and outlook

In this paper we computed twisted equivariant K-groups, as they arise in the context of Kazama-Suzuki models. In the case of Grassmannian coset models the charges obtained from the K-theory match with the conformal field theory charges, as obtained from even (or odd) A-type Cardy states. The K-theory is obtained as a quotient of the representation ring of the denominator group. In particular, the charge lattice satisfies the level-rank duality of the bulk theory. Unlike the WZW models, where the twisted K-theory for $SU(n+1)$, $n \geq 3$, is strictly larger than the charge lattice of Cardy branes, in the coset model case, the Cardy brane charge lattice (at least for the Grassmannian cosets) is precisely given by the twisted equivariant K-theory. However, as we pointed out, there is a subtlety regarding the comparison to D-brane charges in string theory, which seems to be rooted in the GSO-projection.

It would be an interesting check to compute the CFT intersection index in the case of the generalized parafermions, but we shall postpone this to elsewhere. In view of the issue related to the GSO-projection, it would be interesting to extend the above K-theoretical computations to *e.g.*, Gepner models, where an unambiguous string theory description is at hand, and to compare them to the charges computed in [13,18]. More generally, the FHT theorem may be of help in order to shed some more light on the relation between BCFT data and twisted K-theory.

Acknowledgments

It is a great pleasure to thank Constantin Teleman, for many illuminating discussions on K-theory, as well as Matthias Gaberdiel and Greg Moore. I am also very grateful to Volker Braun, Stefan Fredenhagen, Peter Goddard, Andreas Recknagel, and Daniel Roggenkamp for discussions at various stages of this project. I thank St John's College, Cambridge, for a Jenkins Scholarship and acknowledge partial support by EEC contract HPRN-2000-00122.

References

- [1] D.S. Freed, *The Verlinde algebra is twisted equivariant K-theory*, Turkish J. Math. **25** no. 1, 159 (2001); [math.RT/0101038](#).
- [2] D.S. Freed, M.J. Hopkins, C. Teleman, *Twisted K-theory and the Verlinde algebra*, in preparation.
- [3] D.S. Freed, M.J. Hopkins, C. Teleman, *Twisted equivariant K-theory with complex coefficients*, [math.AT/0206257](#).
- [4] R. Minasian and G. W. Moore, *K-theory and Ramond-Ramond charge*, JHEP **9711**, 002 (1997); [hep-th/9710230](#).
- [5] E. Witten, *D-branes and K-theory*, JHEP **9812**, 019 (1998); [hep-th/9810188](#).
- [6] S. Fredenhagen, V. Schomerus, *Branes on group manifolds, gluon condensates, and twisted K-theory*, JHEP **0104**, 007 (2001); [hep-th/0012164](#).
- [7] J. Maldacena, G. Moore, N. Seiberg, *D-brane instantons and K-theory charges*; [hep-th/0108100](#).
- [8] V. Braun, *Twisted K-Theory of Lie Groups*; [hep-th/0305178](#).
- [9] P. Goddard, A. Kent, D. Olive, *Unitary Representations Of The Virasoro And Super-virasoro Algebras*, Commun. Math. Phys. **103**, 105 (1986).
- [10] Y. Kazama, H. Suzuki, *New N=2 Superconformal Field Theories And Superstring Compactification*, Nucl. Phys. B **321**, 232 (1989).
- [11] A. B. Zamolodchikov, V. A. Fateev, *Operator Algebra And Correlation Functions In The Two-Dimensional Wess-Zumino SU(2) X SU(2) Chiral Model*, Sov. J. Nucl. Phys. **43**, 657 (1986)
- [12] D. Gepner, *Field Identification In Coset Conformal Field Theories*, Phys. Lett. B **222**, 207 (1989).
- [13] K. Hori, A. Iqbal, C. Vafa, *D-branes and mirror symmetry*; [hep-th/0005247](#).
- [14] J. Maldacena, G. Moore, N. Seiberg, *Geometrical interpretation of D-branes in gauged WZW models*, JHEP **0107**, 046 (2001); [hep-th/0105038](#).
- [15] W. Lerche, J. Walcher, *Boundary rings and N = 2 coset models*, Nucl. Phys. B **625**, 97 (2002); [hep-th/0011107](#).
- [16] S. Fredenhagen, *Organizing boundary RG flows*; [hep-th/0301229](#).
- [17] M. R. Douglas, B. Fiol, *D-branes and discrete torsion. II*, [hep-th/9903031](#).
- [18] I. Brunner, M. R. Douglas, A. E. Lawrence, C. Römelsberger, *D-branes on the quintic*, JHEP **0008**, 015 (2000); [hep-th/9906200](#).
- [19] W. Lerche, C. Vafa, N. P. Warner, *Chiral Rings In N=2 Superconformal Theories*, Nucl. Phys. B **324**, 427 (1989).
- [20] V. Kac, *Infinite Dimensional Lie Algebras*, Cambridge University Press, (1990).
- [21] M. A. Walton, *Fusion Rules In Wess-Zumino-Witten Models*, Nucl. Phys. B **340**, 777 (1990).

- [22] S. Fredenhagen and V. Schomerus, *Brane dynamics in CFT backgrounds*; **hep-th/0104043**.
- [23] S. Fredenhagen, V. Schomerus, *D-branes in coset models*, JHEP **0202**, 005 (2002); **hep-th/0111189**.
- [24] G. Segal, *Equivariant K-theory*, Inst. Hautes Études Sci. Publ. Math. **34**, 129 (1968).
- [25] M.F. Atiyah, G.B. Segal, *Equivariant K-theory, and completion*, J. Differential Geometry **3**, 1 (1969).
- [26] J. P. May, *Equivariant Homotopy and Cohomology Theory*, CBMS, Regional Conference Series **91**, AMS, 1996.
- [27] M.F. Atiyah, *K-theory Past and Present*, **math.KT/0012213**
- [28] G. Moore, *Lectures on Branes, K-Theory and RR Charges*, Clay Mathematical Institute Lectures; <http://www.physics.rutgers.edu/~gmoore/clay.html>.
- [29] D. S. Freed, *Twisted K-theory and loop groups*; **math.at/0206237**.
- [30] N.E. Steenrod, *Cohomology operations*, Annals of Mathematics Studies **50**, Princeton University Press (1962).
- [31] M. R. Gaberdiel, *Lectures on non-BPS Dirichlet branes*, Class. Quant. Grav. **17**, 3483 (2000); **hep-th/0005029**.
- [32] V. G. Kac, I. T. Todorov, *Superconformal Current Algebras And Their Unitary Representations*, Commun. Math. Phys. **102**, 337 (1985).
- [33] S. G. Naculich, H. J. Schnitzer, *Superconformal coset equivalence from level-rank duality*, Nucl. Phys. B **505**, 727 (1997); **hep-th/9705149**.
- [34] L. Hodgkin, *The equivariant Künneth theorem in K-theory*, in *Topics in K-theory. Two independent contributions*, Lecture Notes in Math. **496**, Springer, Berlin, 1975.
- [35] V.P. Snaith, *On the Künneth formula spectral sequence in equivariant K-theory*, Proc. Cambridge Philos. Soc. **72**, 167, (1972).
- [36] H.V. Pittie, *Homogeneous vector bundles on homogeneous spaces*, Topology **11**, 199 (1972).
- [37] D. Gepner, *Fusion Rings and Geometry, Fusion Rings And Geometry*, Commun. Math. Phys. **141**, 381 (1991).
- [38] A. Pressley, G. Segal, *Loop Groups*, Oxford, 1986.
- [39] R. Hartshorne, *Algebraic Geometry*, Springer, 1977.
- [40] D.S. Freed, M.J. Hopkins, C. Teleman, *Twisted K theory and Loop group representations*, to appear.
- [41] J.F. Adams, *Vector Fields on Spheres*, Ann. of Math. **74**, 603 (1962).